

Were these symptoms present on most days for at least one month?*
Kentucky BRFSS 2001

Geographic Groups	Total Respond.#	Yes			No		
		N	%	C.I.(95%)	N	%	C.I.(95%)
TOTAL	3917	2607	63.7	(61.4-65.9)	1310	36.3	(34.1-38.6)
ADDs							
Barren River	277	189	65.5	(58.6-71.8)	88	34.5	(28.2-41.4)
Big Sandy	268	190	69.4	(62.8-75.2)	78	30.6	(24.8-37.2)
Bluegrass	251	163	66.3	(59.2-72.7)	88	33.7	(27.3-40.8)
Buffalo Trace	199	134	69.6	(61.1-76.9)	65	30.4	(23.1-38.9)
Cumberland Valley	267	181	65.6	(58.7-71.8)	86	34.4	(28.2-41.3)
FIVCO	334	225	66.0	(59.9-71.6)	109	34.0	(28.4-40.1)
Gateway	298	196	64.2	(57.6-70.3)	102	35.8	(29.7-42.4)
Green River	204	140	64.9	(56.7-72.2)	64	35.1	(27.8-43.3)
Kentucky River	283	213	70.2	(63.4-76.2)	70	29.8	(23.8-36.6)
KIPDA	257	160	59.5	(52.3-66.2)	97	40.5	(33.8-47.7)
Lake Cumberland	266	170	61.5	(54.4-68.1)	96	38.5	(31.9-45.6)
Lincoln Trail	249	154	61.0	(53.8-67.7)	95	39.0	(32.3-46.2)
Northern Kentucky	267	162	59.3	(52.4-65.9)	105	40.7	(34.1-47.6)
Pennyrile	219	147	64.1	(56.3-71.3)	72	35.9	(28.7-43.7)
Purchase	278	183	65.3	(58.8-71.2)	95	34.7	(28.8-41.2)

*This question is asked to respondents who reported having pain, aching, stiffness or swelling in or around a joint.

#Use caution in interpreting data for total respondents less than 50. N = Cell Size, NH = Non-Hispanic, %

= Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level).

Percentages are weighted to population characteristics.

Unless stated differently above, Missing, 'Don't know', and Refused responses are excluded.